

Reaction plane dependence of neutral pion production in center-of-mass energy of 200 GeV Au+Au collisions at RHIC-PHENIX

Yoki Aramaki, for the PHENIX Collaboration

Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

Abstract

The integrated luminosity of RHIC-Year 2007 Au+Au run is $813 \mu b^{-1}$, which is 3.5 times larger than that in RHIC-Year 2004 Au+Au run. Additionally, a new detector was installed to determine reaction plane more precisely. This detector is expected to provide better resolution for reaction plane determination. These advantages enable us to precisely measure the path length dependence of π^0 suppression and discuss parton energy loss mechanism more thoroughly. We report the recent results for the reaction plane dependence of π^0 production.

Key words: Parton energy loss, Collective flow

PACS: 25.75.-q, 25.75.Ld, 21.65.Qr

1. Introduction

It has been observed in central Au+Au collisions at Relativistic Heavy Ion Collider (RHIC) that the yield of neutral pions at high transverse momentum ($p_T > 5$ GeV/ c) is strongly suppressed compared to the one expected from p+p collisions scaled by the number of binary collisions. This suppression is considered to be due to the energy lost by hard scattered partons in the medium (jet quenching), which results in a decrease of the yield at a given p_T . Many theoretical models have been proposed to understand the parton energy loss mechanism. For one thing, GLV method[1] as one of the calculations predicts that the magnitude of energy loss is proportional to the path length if the medium has static density. Studying the path length dependence of energy loss should help the understanding of energy loss process.

Email address: aramaki@cns.s.u-tokyo.ac.jp (Yoki Aramaki, for the PHENIX Collaboration).

2. Azimuthal angle dependence of R_{AA}

Recently theoretical models (ASW[2], HT[3] and AMY[4]) to describe parton energy loss mechanism which involve the time-evolution of the medium produced at RHIC have been proposed. These models succeeded in reproducing the centrality dependence of $R_{AA}(p_T)$. These theoretical curves and the preliminary data from PHENIX is shown in Fig. 1[5]. The central Fig. 1 shows expected curves from these models.

As shown on the right panel of Fig. 1, these models are still unable to reproduce the azimuthal angle dependence of R_{AA} , even if we compare the curve with the largest variation, the $p_T = 15$ GeV/c of the ASW model to data at much lower p_T ($5 < p_T < 8$ GeV/c).




Fig. 1. Left: R_{AA} in Au+Au collisions at 0-5 % (top) and 20-30 % (bottom) centrality calculated in the ASW, HT and AMY models compared with data from PHENIX[6]. Central: R_{AA} as a function of azimuthal angle at $p_T = 10$ GeV/c (solid line) and $p_T = 15$ GeV/c (dashed line) for all three models at 20-30 % centrality. Right: R_{AA} as a function of azimuthal angle ($5 < p_T < 8$ GeV/c) at centrality 20-30% and blue dashed line is ASW model at $p_T = 15$ GeV/c.

When partons lose energy in the medium, the ratio of $R_{AA}(\Delta\phi = 0)$ to $R_{AA}(\Delta\phi = \pi/2)$ should be different for each centrality bin. Fig. 2 shows the azimuthal angle dependence of R_{AA} at $3 < p_T < 5$ GeV/c. As shown in Fig. 2, $R_{AA}(3 < p_T < 5$ GeV/c, $\Delta\phi = 0)$ is about two times larger than $R_{AA}(3 < p_T < 5$ GeV/c, $\Delta\phi = \pi/2)$ for all centrality bins. This may be in part be an effect of collective flow. Since the influence of collective flow at high p_T should be small, we need to measure $R_{AA}(p_T, \Delta\phi)$ at higher p_T to minimize its effect.

3. $v_2(\pi^0)$ at high p_T

Reaction plane detector (RxnP) was installed in RHIC-Year 2007 and reaction plane determination accuracy has been improved by a factor of two as compared to that in RHIC-Year 2004. Fig. 3 shows $v_2(\pi^0)$ as a function of p_T for each 20 % centrality bin. This detector enabled us to measure $v_2(\pi^0)$ up to 14 GeV/c. Measured $v_2(\pi^0)$ are non-zero for all centrality bins. Additionally, two assumed function is fitted to this data. One is linear function ($f(v_2) = a \cdot p_T + b$) and the other is constant value ($f(v_2) = c$). At centrality 0-20 %, values of χ^2/NDF for constant and linear function are 4.45/5 and 4.34/4, respectively. At centrality 20-40 %, values of χ^2/NDF for




Fig. 2. Azimuthal angle dependence of R_{AA} integrated over $3 < p_T < 5$ GeV/c[7].

constant and linear function are 4.39/5 and 1.49/4, respectively. At centrality 40-60 %, values of χ^2/NDF for constant and linear function are 2.23/5 and 2.21/4, respectively. These results indicate that the values of $v_2(\pi^0)$ in most central and peripheral collisions tend to be constant while in mid-central collisions tend to decrease.




Fig. 3. $v_2(\pi^0)$ as a function of p_T for each 20 % centrality bin. Red and blue lines show constant and linear function, respectively. Red and blue lines are fitted to data from 6 GeV/c.

4. Summary and outlook

Study of path length and azimuthal angle dependence of R_{AA} with the new reaction plane detector has been started. The nuclear modification factor, $R_{AA}(5 < p_T < 8 \text{ GeV}/c)$ as a function of azimuthal angle has been compared to theoretical models. Even though the models are in good agreement with the (azimuthally integrated) R_{AA} , so far they could not reproduce the measured azimuthal angle dependence. The measurement of $v_2(\pi^0)$ has now been extended to 14 GeV/c. For the most central (0-20%) and peripheral (40-60%) collisions, the elliptic flow tends to be constant (instead of decreasing monotonically with p_T), while for mid-central collisions (20-40%) we observe a decrease with p_T .

With the new data we will be able to measure the dependence of R_{AA} on azimuthal angle up to higher transverse momenta than ever before. We can also estimate the path length by measuring the azimuthal angle from reaction plane and mapping it into the shape of the participant region, which can be calculated by Glauber model for each impact parameter (centrality).

References

- [1] M. Gyulassy, P. Levai and I.Vitev Phys. Lett. B**538**, 282 (2002).
- [2] T. Renk *et al.* Phys. Rev. C**75** 031902 (2007).
- [3] A. Majumder *et al.* Phys. Rev. C**76** 041902 (2007).
- [4] G-Y. Qin *et al.* Phys. Rev. C**76** 064907 (2007).
- [5] S.A. Bass *et al.* J. Phys. G**35** 104064 (2008).
- [6] M. Shimomura *et al.* Nucl. Phys. A **774** 457-460, (2006).

- [7] S.S. Adler *et al* (*PHENIX*). *Phys. Rev. C* **76** 034904, (2007).